

# Temperature-mediated plasticity and genetic differentiation in egg size and hatching size among populations of *Crepidula* (Gastropoda: Calyptraeidae)

RACHEL COLLIN\* and MARICELA Z. SALAZAR

*Smithsonian Tropical Research Institute, Box 0843-03092, Balboa, Republic of Panama*

Received 7 August 2009; accepted for publication 29 September 2009

Offspring size is a key characteristic in life histories, reflecting maternal investment per offspring and, in marine invertebrates, being linked to mode of development. Few studies have focused explicitly on intraspecific variation and plasticity in developmental characteristics such as egg size and hatching size in marine invertebrates. We measured over 1000 eggs and hatchlings of the marine gastropods *Crepidula atrasolea* and *Crepidula ustulatulina* from two sites in Florida. A common-garden experiment showed that egg size and hatching size were larger at 23 °C than at 28 °C in both species. In *C. ustulatulina*, the species with significant genetic population structure in cytochrome oxidase I (COI), there was a significant effect of population: Eggs and hatchlings from the Atlantic population were smaller than those from the Gulf. The two populations also differed significantly in hatchling shape. Population effects were not significant in *C. atrasolea*, the species with little genetic population structure in COI, and were apparent through their marginal interaction with temperature. In both species, 60–65% of the variation in egg size and hatching size was a result of variation among females and, in both species, the population from the Atlantic coast showed greater temperature-mediated plasticity than the population from the Gulf. These results demonstrate that genetic differentiation among populations, plastic responses to variation in environmental temperature, and differences between females all contribute significantly to intraspecific variation in egg size and hatching size. © 2010 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2010, **99**, 489–499.

ADDITIONAL KEYWORDS: evolution of development – maternal effects – offspring size – phylogeography.

## INTRODUCTION

Documenting patterns of intraspecific variation is vital to understanding the evolutionary dynamics of any trait (Bernardo, 1996). Despite this, studies of intraspecific variation are uncommon in certain fields. Several reviews have called specifically for more detailed studies of intraspecific variation in offspring size or propagule size (Bernardo, 1996; Marshall & Keough, 2008). Bernardo (1996) argues that the commonly studied patterns in interspecific variation do not provide adequate information to understand the evolutionary dynamics of offspring size, and that

studies of within and among population variation are vital to make progress in this area. He specifically cites the traditional focus on the evolution of optimal offspring size, instead of on the evolution of reaction norms in offspring size or other aspects of plasticity, as an impediment to understanding evolutionary patterns in offspring size. Studies of intraspecific variation in propagule size in marine invertebrates are particularly rare (Marshall & Keough, 2008; Moran & McAlister, 2009) and plasticities are seldom examined.

## PROPAGULE SIZE IN MARINE INVERTEBRATES

Egg size and hatching size have been the focus of many studies of life histories of marine invertebrates, yet there are few explicit studies of their intraspecific variation (Hadfield & Strathmann, 1996; Jones, Todd

\*Corresponding author. Smithsonian Tropical Research Institute, MRC 0580-08, Unit 9100 Box 0948, APO AA 34002, USA. E-mail: collinr@si.edu

& Lambert, 1996; Marshall & Keough, 2008). Interspecific differences in these characteristics are often extreme and are associated with different modes of development: planktonic versus benthic, feeding versus nonfeeding, and direct versus indirect. Interspecific variation in egg size and/or hatching size can also affect fertilization rate and success (Levitan, 2006), development time (Kohn & Perron, 1994; Emler, 1995), predation rates by zooplankton (Allen, 2008), hatching size (Kohn & Perron, 1994), settlement size (George, Cellario & Fenaux, 1990; Hart, 1995; Allen, Zakas & Podolsky, 2006), and juvenile growth rate and survival (Rivest, 1983; Moran & Emler, 2001; Marshall & Keough, 2003). Intraspecific variation in offspring size generally has similar but smaller effects than intraspecific variation on many stages of the life cycle.

The existing literature suggests that there is widespread intraspecific variation in egg size and hatching size (Hadfield & Strathmann, 1996; Marshall & Keough, 2008), although there is no clear picture of how this variation is partitioned within and among broods, females, and populations. Similarly, few studies have examined how plastic responses to environmental factors such as temperature, or maternal effects such as responses to maternal size or nutrition effect offspring size in marine invertebrates.

The studies that do examine intraspecific variation in offspring size have found relationships between egg size or hatching size, and maternal size (Ito, 1997; Chaparro *et al.*, 1999; Marshall, Styan & Keough, 2000), temperature (Skadsheim, 1989; Honkoop & Van Der Meer, 1998), and latitude or tidal height (Hadfield, 1989; Lardies & Castilla, 2001). In addition, egg sizes have been found to vary among populations (Barnes & Barnes, 1965; Jones *et al.*, 1996), and can show patterns consistent with seasonal or lunar periodicity (Lessios, 1987).

#### INTRASPECIFIC VARIATION IN ATLANTIC AND GULF *CREPIDULA*

We used a common garden experiment to compare Atlantic and Gulf populations of two species of *Crepidula* (Gastropoda: Calyptraeidae), to document temperature-mediated plasticity in egg size and hatching size, and to examine differences among populations. In one species, the populations are reciprocally monophyletic at the cytochrome oxidase I (COI) locus. In the other species, the two populations share haplotypes. Therefore, this experiment also tests the hypothesis that the species with more population structure in COI also shows more differences between populations in life history traits. If divergence in COI reflects levels of divergence across the entire genome, as is often assumed, then divergence

in COI should be associated with divergences in other traits, such as offspring size.

Development and reproduction are well studied in calyptraeid gastropods and phylogenetic studies have demonstrated rapid evolution of mode of development and concomitant changes in egg size (Collin, 2004; Collin *et al.*, 2007). *Crepidula atrasolea* Collin 2000 and *Crepidula ustulatulina* Collin 2002 (the 'southern species' of *C. convexa*; Collin, 2001) co-occur along the coast of Florida and both brood, have large eggs, and previously reported variation in egg size. *Crepidula atrasolea* has direct development with embryos that hatch as crawling juveniles. Egg diameter is reported as 335  $\mu\text{m}$  (Collin, 2003) and hatching size as 900  $\mu\text{m}$  (*Crepidula cf. plana*; Hoagland, 1986) or 1002  $\mu\text{m}$  (Collin, 2003). Development of *C. ustulatulina* includes lecithotrophic larvae that swim for minutes to hours prior to settling. Egg diameter is reported as 300–340  $\mu\text{m}$  and hatchling shell lengths are reported as 744  $\mu\text{m}$  (Collin, 2003), 840  $\mu\text{m}$  (*Crepidula cf. convexa*; Hoagland, 1986) or 630–700  $\mu\text{m}$  (*Crepidula cf. convexa*; Hoagland, 1984).

Phylogeographic analysis has shown significant differences in population structure in mitochondrial (mt)COI between the two species (Collin, 2001). *Crepidula atrasolea* shows little genetic structure along the coast of Florida. Populations from Mote Marine Laboratory, near Sarasota on the Gulf coast and Fort Pierce on the Atlantic coast of Florida share COI haplotypes that differ by one or two base pairs and are not reciprocally monophyletic. By contrast *C. ustulatulina* has significant population structure and the reciprocally monophyletic COI haplotypes from Mote and Fort Pierce differ by 2% (Collin, 2001). Animals from both populations interbreed successfully in the laboratory (R. Collin, unpubl. data), supporting the conspecific identity of the different populations.

In the present study, we used animals from these two populations raised at two temperatures to determine: (1) whether egg size and hatching size show temperature-mediated plasticity; (2) whether the species with distinct COI haplotypes among populations showed greater population-level differences in egg size and hatching size than the species that shares COI haplotypes among populations; and (3) whether population  $\times$  temperature effects are also greater in the species with distinct COI haplotypes among populations than the species that shares COI haplotypes.

#### MATERIAL AND METHODS

Brooding females and small juveniles of both species were collected from less than 1 m depth at low tide near the breakwater at the Harbor Branch

Oceanographic Institute in Fort Pierce on the Atlantic coast of Florida, and in the seagrass meadow in Sarasota Bay north of the Mote Marine Laboratory on the Gulf coast of Florida. Pairs of hatchlings from the same population were transferred to 350-mL plastic cups and raised for approximately 1 year in incubators at either 23 °C or 28 °C (within the range of temperatures experienced during the reproductive season). The larger animal in each pair became female after 3–4 months, and the smaller animal developed as male; thus, eggs were obtained from only one animal per pair. Water was changed every other day and  $3.86 \times 10^6$  cells of *Isochrysis galbana* culture was added to the cups daily. At large (reproductive) sizes, this was probably a limiting ration because the water was cleared hours before the next scheduled feeding.

Females were checked twice each day for new eggs, which could be seen through the transparent plastic of the cups without disturbing the females. Uncleaved eggs were collected, placed on a slide with a no. 1 cover slip, supported by large 'clay feet' to ensure that pressure from the cover slip did not squash or alter the egg size or shape in any way, and photographed. We aimed to photograph twenty round, uncleaved eggs from each female, but could not always find 20 when the females were small or cleavage had begun. Hatchlings were collected within 12 h of hatching naturally from the egg capsule, and preserved in 70% ethanol. The shells were stained blue, oriented flat and photographed. Prior to photographing, the eggs or hatchlings from each female, a stage micrometer was photographed with the same magnification. Only a single brood of eggs and a single brood of hatchlings were measured from each female.

Using the software IMAGEJ (Abramoff, Magelhaes & Ram, 2004), the micrometer photograph was used to calibrate the measurements, and the Shape Descriptor plug-in was used to measure the area, diameter or major and minor axes, aspect ratio (major axis/minor axis) and roundness ( $4 \times \text{area} / \pi \times \text{diameter}^2$ ) of each egg or hatchling. Eggs with a roundness of less than 94 were eliminated from the analysis because eggs elongate prior to cleavage and changes in shape can alter size estimates.

Data were analysed using the REML (restricted maximum likelihood) algorithm analysis of variance as implemented in JMP, version 5.1 (SAS Institute Inc.) with eggs produced by each female nested within population as a random effect. REML can provide more reliable estimates of variance for unbalanced models than standard analysis of variance (Quinn & Keough, 2002) and is recommended for analysing random effects. *P*-values for the random nested effect were calculated on the likelihood value of the model using a *Z*-test and Tukey's honestly significant differ-

ence (HSD) test were used to compare group means. Egg diameter was used since it is the most commonly used measure of egg size in the literature. Because there were significant differences between populations and temperatures in hatchling shapes, hatchling area was used as the measure of hatchling size for statistical analyses. Correlations between female size and egg size or hatching size were examined using analysis of covariance (ANCOVA) on average egg size for each brood, with population and temperature as direct effects and female size at the time the brood was laid as a covariate. Stepwise removal was used to eliminate nonsignificant interactions. These datasets were smaller than the total egg and hatchling datasets because female size was not measured for all of the animals when eggs and hatchlings were collected.

Variation in egg size is usually inferred to reflect variation in energy content and therefore maternal investment. However, egg size does not always reflect energy content because levels of hydration as well as allocation to different yolk components can vary among eggs and among mothers (McEdward & Carson, 1987; McEdward & Coulter, 1987; Moran & McAlister, 2009). It is not easy to verify the relationship between size and energy content of individual eggs because it is difficult to obtain accurate measures of energy content for such tiny samples. If egg size does reflect energy content, then egg size should be correlated with hatching size. Unfortunately, removing eggs from brooding *Crepidula* results in the abandonment of the entire brood and the eggs do not develop normally away from their mother. Therefore, we examined the correlation between egg size and hatching size from different broods from the same mother using ordinary least squares regression on brood means.

## RESULTS

### EGG SIZE

The overall average egg diameter was 339 µm for *C. atrasolea* and 304 µm for *C. ustulatulina* (Table 1). For both species, smaller sample sizes were obtained in the high temperature treatment (Table 1). This was primarily because eggs cleaved quickly at high temperatures, making it difficult to collect eggs when they were still round. In addition, adult snails in the high temperature treatment did not appear to produce broods as frequently as those at the lower temperature.

For *C. atrasolea*, there was a significant effect of temperature and female on egg size but no interaction between temperature and population (Fig. 1, Tables 1, 2) and no direct effect of population. Egg size was smaller (overall average 333 µm) at 28 °C than in the 23 °C treatment (overall average 341 µm). Eggs

**Table 1.** Summary of egg size and hatching size in *Crepidula atrasolea* and *C. ustulatulina*

|                               | Fort Pierce<br>(23 °C) | Fort Pierce<br>(28 °C) | Mote<br>(23 °C) | Mote<br>(28 °C) | Total | Individual females<br>mean (range) |
|-------------------------------|------------------------|------------------------|-----------------|-----------------|-------|------------------------------------|
| <i>Crepidula atrasolea</i>    |                        |                        |                 |                 |       |                                    |
| Egg diameter                  |                        |                        |                 |                 |       |                                    |
| Average (µm)                  | 344.4                  | 330.1                  | 337.7           | 337.1           | 338.8 | 299–394                            |
| SD (µm)                       | 23.2                   | 19.6                   | 16.4            | 17.2            | 20.6  | 9.2 (3.3–55.7)                     |
| Number of eggs                | 556                    | 240                    | 645             | 214             | 1655  | 18 (10–23)                         |
| Number of females             | 30                     | 14                     | 33              | 12              | 89    | 89                                 |
| Hatching length               |                        |                        |                 |                 |       |                                    |
| Average (µm)                  | 1074                   | 1005                   | 1025            | 987             | 1026  | 850–1227                           |
| SD (µm)                       | 93.2                   | 93.4                   | 97.1            | 103.7           | 102.0 | 57.2 (28.0–141.6)                  |
| Number of eggs                | 511                    | 322                    | 663             | 404             | 1880  | 17.7 (6–20)                        |
| Number of females             | 27                     | 20                     | 35              | 25              | 107   | 107                                |
| <i>Crepidula ustulatulina</i> |                        |                        |                 |                 |       |                                    |
| Egg diameter                  |                        |                        |                 |                 |       |                                    |
| Average (µm)                  | 306.3                  | 281.5                  | 307.5           | 301.5           | 303.5 | 248–354                            |
| SD (µm)                       | 25.4                   | 22.0                   | 17.5            | 17.7            | 22.0  | 10.0 (3.8–61.0)                    |
| Number of eggs                | 376                    | 133                    | 568             | 224             | 1301  | 17.6 (9–20)                        |
| Number of females             | 22                     | 8                      | 30              | 13              | 73    | 73                                 |
| Hatching length               |                        |                        |                 |                 |       |                                    |
| Average (µm)                  | 932.1                  | 873.4                  | 930.7           | 905.8           | 916   | 740.0–1160.5                       |
| SD (µm)                       | 89.3                   | 95.6                   | 88.3            | 101.2           | 95.5  | 53.7 (19.7–133.9)                  |
| Number of eggs                | 578                    | 287                    | 645             | 513             | 2023  | 18.9 (9–20)                        |
| Number of females             | 30                     | 16                     | 34              | 28              | 108   | 108                                |

**Table 2.** Restricted maximum likelihood analysis of variance of egg size of *Crepidula atrasolea* and *Crepidula ustulatulina*

|                               |      |                |  |         |                    |
|-------------------------------|------|----------------|--|---------|--------------------|
| <i>Crepidula atrasolea</i>    |      |                |  |         |                    |
| Fixed effects                 | d.f. | SS             |  | F-ratio | P-value            |
| Temperature                   | 1    | 613.47         |  | 4.49    | <b>0.03</b>        |
| Population                    | 1    | 0.09           |  | 0.0007  | 0.98               |
| Temperature × population      | 1    | 412.17         |  | 3.02    | 0.08               |
| Random effect                 | d.f. | Variance ratio |  | Z-ratio | P-value            |
| Female [population]           | 88   | 2.00           |  | 6.20    | <b>&lt; 0.0001</b> |
| <i>Crepidula ustulatulina</i> |      |                |  |         |                    |
| Fixed effects                 | d.f. | SS             |  | F-ratio | P-value            |
| Temperature                   | 1    | 1691.06        |  | 10.36   | <b>0.001</b>       |
| Population                    | 1    | 865.53         |  | 5.30    | <b>0.02</b>        |
| Temperature × population      | 1    | 605.89         |  | 3.71    | <b>0.05</b>        |
| Random effect                 | d.f. | Variance ratio |  | Z-ratio | P-value            |
| Female (population) random    | 70   | 2.00           |  | 5.50    | <b>&lt; 0.0001</b> |


Significant *P*-values in bold.

from 28 °C and 23 °C from Fort Pierce differ from each other but not from the eggs from Mote (Tukey's HSD test; Fig. 1). Most of the variation in egg size (65%) was a result of the significant effects of individual females.

For *C. ustulatulina*, there were significant effects of temperature, population, and female on egg diameter (Fig. 1, Tables 1, 2), and a marginally significant

interaction between temperature and population (Table 2). Egg size was larger (overall average 307 µm) at 23 °C than it was at 28 °C (overall average 294 µm). Again, the majority of variation (62%) in egg size was a result of the effect of female.

There was no clear effect of maternal size on egg size in either species. For *C. ustulatulina*, there was



**Figure 1.** Effects of rearing temperature and population of origin on egg size and hatching size in *Crepidula atrasolea* and *Crepidula ustulatulina*. Restricted maximum likelihood analysis of variance showed a significant effect of temperature in all four cases and a significant effect of population for egg size and hatching size in *C. ustulatulina*. Error bars indicate the standard errors.

no significant effect of maternal size on egg size (ANCOVA:  $F = 0.02$ ;  $N = 38$ ;  $P > 0.9$ ) in the total dataset, nor when the dataset was partitioned by population or temperature. There was no significant effect of maternal shell length on egg diameter in *C. atrasolea* when eggs from both temperature treatments were analysed together (ANCOVA:  $F = 0.25$ ;  $N = 77$ ;  $P > 0.6$ ). There was also no significant effect of temperature for this dataset, which is smaller than our total dataset, although there was a significant interaction between the effects of temperature and maternal shell length ( $P < 0.003$ ;  $N = 77$ ). At 23 °C, egg size decreases with maternal size ( $r^2 = 0.12$ ;  $N = 59$ ,  $P = 0.007$ ) but, at 28 °C, egg size increases with maternal size ( $r^2 = 0.24$ ;  $N = 18$ ;  $P = 0.04$ ).

#### HATCHLING SIZE

The average hatching length was 1026  $\mu\text{m}$  for *C. atrasolea* and 916  $\mu\text{m}$  for *C. ustulatulina* (Table 1). For *C. atrasolea*, there was a significant effect of temperature and female on hatching area, and no significant effect of population or temperature  $\times$  population (Fig. 1, Tables 1, 3). The average hatching length was smaller (995  $\mu\text{m}$ ) at 28 °C than at 23 °C (1046  $\mu\text{m}$ ).

Again, most of the variation in hatching size (65%) was a result of the significant effect of female.

For *C. ustulatulina*, there was a significant effect of temperature, population, and female on hatching area (Fig. 1, Tables 1, 3), and no significant temperature  $\times$  population effect. Hatching length was larger (931  $\mu\text{m}$ ) at 23 °C than it was at 28 °C (894  $\mu\text{m}$ ). Hatchlings from Fort Pierce were smaller than those from Mote at both temperatures. Again, the majority of variation (65%) in hatching size was as result of the effect of female.

The effect of female size explained very little of the variation in hatching size. Hatching area was not significantly correlated with maternal size for *C. atrasolea* for the total dataset (ANCOVA:  $F = 0.55$ ;  $N = 107$ ;  $P > 0.7$ ), nor when the dataset was partitioned by temperature or population. Hatching size showed no effect of maternal size in *C. ustulatulina* for the entire dataset (ANCOVA:  $F = 2.62$ ;  $N = 101$ ;  $P > 0.1$ ). However, when the dataset was partitioned by population, the Mote population showed a small but significant increase in hatching size with maternal shell length ( $N = 58$ ;  $r^2 = 0.10$ ;  $P < 0.05$ ). The Fort Pierce population did not show any significant relationship between the two variables.

**Table 3.** Restricted maximum likelihood analysis of variance of hatching size of *Crepidula atrasolea* and *Crepidula ustulatulina*

| <i>Crepidula atrasolea</i>      |      |                       |         |                 |
|---------------------------------|------|-----------------------|---------|-----------------|
| Fixed effects                   | d.f. | SS                    | F-ratio | P-value         |
| Temperature                     | 1    | $4.14 \times 10^{10}$ | 7.45    | <b>0.006</b>    |
| Population                      | 1    | $1.62 \times 10^{10}$ | 2.92    | 0.09            |
| Temperature $\times$ population | 1    | $7.88 \times 10^9$    | 1.42    | 0.23*           |
| Random effect                   | d.f. | Variance ratio        | Z-ratio | P-value         |
| Female (population) random      | 104  | 1.81                  | 6.70    | < <b>0.0001</b> |
| <i>Crepidula ustulatulina</i>   |      |                       |         |                 |
| Fixed effects                   | d.f. | SS                    | F-ratio | P-value         |
| Temperature                     | 1    | $1.67 \times 10^{10}$ | 4.97    | <b>0.03</b>     |
| Population                      | 1    | $1.72 \times 10^{10}$ | 5.13    | <b>0.03</b>     |
| Temperature $\times$ population | 1    | $5.61 \times 10^9$    | 1.67    | 0.20            |
| Random effect                   | d.f. | Variance ratio        | Z-ratio | P-value         |
| Female (population) random      | 105  | 1.92                  | 6.81    | < <b>0.0001</b> |

\*Removal of the nonsignificant interaction effect does not alter the result. Significant *P*-values in bold.

There is some consistency in relative offspring size from brood to brood, suggesting a maternal component. Hatchling size in one brood was positively correlated with egg diameter in a different brood from the same female, although this explained only a small part of the variance in hatchling size (*C. atrasolea*:  $r^2 = 0.11$ ;  $N = 39$ ;  $P = 0.04$ ; *C. ustulatulina*  $r^2 = 0.11$ ;  $N = 56$ ;  $P = 0.01$ ; Fig. 2).

#### HATCHLING SHAPE

Analysis of hatchling shape was based on aspect ratio, the ratio of major to minor axes. For *C. atrasolea*, there was a significant effect of temperature and female, but not of population (Table 4) on aspect ratio. Hatchlings at 23 °C were longer and thinner than those in the high temperature treatment (Fig. 3). A substantial amount of the variation in shape (40%) was a result of differences among females.

For *C. ustulatulina*, temperature, population, and female were significant effects on hatchling shape (Table 3). Similar to *C. atrasolea*, hatchlings at low temperature were longer and thinner than the ones in the high temperature treatment (Fig. 3). In addition, the two populations differed in hatchling shape: hatchlings from Fort Pierce are significantly longer and thinner than those from Mote. Thirty-five percent of the variation was a result of differences among females.


#### DISCUSSION

The data obtained in the present study on egg size and hatching size in these two species of *Crepidula* show that temperature-mediated plasticity, genetic differentiation among populations, and differences between females, all contribute to significant variation in off-

spring size and shape. In both species, the majority of variation in egg size and hatching size was a result of differences among females. Similarly, in both species, egg size and hatching size decreased significantly with increasing temperature, and this pattern was more pronounced in the Atlantic population than the Gulf population. The species differed in the effect of population on offspring size: In *C. ustulatulina*, the species that shows populations differentiation in COI, there was a strongly significant difference in egg size and hatching shape between populations and a significant difference in hatchling size. In *C. atrasolea*, the species lacking differentiation in COI, there was no evidence of differences among populations in egg size, hatching size, or hatching shape. Finally, although there was a significant temperature–population interaction only for egg size, the animals from Fort Pierce showed a pattern of greater response to temperature than those from Mote for all three variables. The possible causes and consequences of these different results are discussed below.

#### ARE THE EFFECTS OF TEMPERATURE ADAPTIVE?


Egg size and hatching size both show a phenotypic response to temperature, with larger offspring being produced at the lower temperature in both species. Such decreases in egg size and hatching size with increasing temperature are common not just in marine invertebrates (Honkoop & Van Der Meer, 1998), but also in insects (Ernsting & Isaaks, 2000; Fischer, Brakefield & Zwaan, 2003) and marine fishes (Bengtson, Barkman & Berry, 1987; Kokita, 2003). Many studies interpret such temperature-mediated plasticity as an adaptive response linked to differential juvenile growth or survival at different temperatures, and seek to quantify reproductive output under



**Figure 2.** Scatter plots of average egg size for a single brood for each female versus average hatching size for a single brood for the same female. A, *Crepidula atrasolea* (trend line is the ordinary least squares (OLS) regression;  $r^2 = 0.11$ ;  $N = 39$ ;  $P = 0.037$ ). B, *Crepidula ustulatulina* (trend line is OLS;  $r^2 = 0.11$ ;  $N = 56$ ;  $P = 0.013$ ).

different conditions (Fischer *et al.*, 2003). Other studies have interpreted this response as a physiological response to stress (George, 1996), either directly from elevated temperatures or via nutritional stress as a result of a higher metabolic rate at higher temperatures. Finally, Van Voorhies (1996) suggested that temperature effects are ubiquitous on cell size in ectotherms.

Stress appears unlikely to be operating in the case of *Crepidula* offspring size. The temperatures employed in the present study were not physiologically stressful for adult animals and were typical of temperatures experienced in the field. Water temperatures at the collection sites for both populations showed similar annual ranges and temperatures in considerable excess of 28 °C. Water temperature mea-



**Figure 3.** Effects of rearing temperature and population of origin on hatchling shape for *Crepidula atrasolea* and *Crepidula ustulatulina*. Restricted maximum likelihood analysis of variance showed a significant effect of temperature on both species as well as a significant effect of population on *C. ustulatulina*. Error bars indicate the standard errors.

sured at mid-day once a month for 5 years from the collecting site in Fort Pierce ranged from 13.8–37.7 °C, with an average of 25.5 °C. Four months of the year have average temperatures higher than 28 °C (S. Reed, pers. comm.). Ten years of monthly data from the seagrass meadow near Mote Marine Laboratory are similar and ranged from 12.8–37.2 °C, with an average of 24.4 °C. The 75% quartile was 28.91 °C (K. Dixon, pers. comm.).

Stress could also have resulted from food limitation; however, it is unclear in which direction nutritional stress would be expected to act. Nutritional stress could have the direct consequence of reduced egg size as a result of reduced resources available to the female (Fox, 1993; Bertram & Strathmann, 1998). However, it could also induce an adaptive maternal response: Because juveniles of both species are unlikely to disperse far from the mother, mothers might be expected to produce larger propagules under nutritional stress (Qian, 1994) to increase their chances of survival. The first scenario suggests that if higher temperatures or larger sizes result in nutritional stress, egg size should decrease with tempera-

**Table 4.** Restricted maximum likelihood analysis of variance of hatchling shape as measured by aspect ratio

| <i>Crepidula atrasolea</i>      |      |                       |         |                    |
|---------------------------------|------|-----------------------|---------|--------------------|
| Fixed effects                   | d.f. | SS                    | F-ratio | P-value            |
| Temperature                     | 1    | 0.0048                | 4.3     | <b>0.038</b>       |
| Population                      | 1    | $1.25 \times 10^{-7}$ | 0.0001  | 0.99               |
| Temperature $\times$ population | 1    | $3.7 \times 10^{-5}$  | 0.033   | 0.86               |
| Random effect                   | d.f. | Variance ratio        | Z-ratio | P-value            |
| Female (population) random      | 104  | 0.66                  | 6.40    | <b>&lt; 0.0001</b> |
| <i>Crepidula ustulatulina</i>   |      |                       |         |                    |
| Fixed effects                   | d.f. | SS                    | F-ratio | P-value            |
| Temperature                     | 1    | 0.011                 | 4.90    | <b>0.027</b>       |
| Population                      | 1    | 0.11                  | 46.73   | <b>&lt; 0.0001</b> |
| Temperature $\times$ population | 1    | 0.00004               | 0.019   | 0.89               |
| Random effect                   | d.f. | Variance ratio        | Z-ratio | P-value            |
| Female (population) random      | 105  | 0.53                  | 6.31    | <b>&lt; 0.0001</b> |

Significant *P*-values in bold.

ture and with maternal size, consistent with our results indicating that egg size decreases with temperature. The second scenario suggests that egg size should increase with nutritional stress and therefore with temperature and with size. This is consistent with our result that egg size increased with maternal size, for *C. atrasolea* at 28 °C, and in *C. ustulatulina* from Mote. Direct tests of the effects of food limitation must be made to determine exactly how this factor contributes to variation in egg size in *Crepidula*.

It is difficult to address the two other possible explanations: adaptive trade-offs and physiological allometry. We did not collect data on the survival or fitness of offspring of different sizes at different temperatures, or the offspring size–clutch size trade-offs necessary to examine the effects of temperature in terms of optimal fitness and survival. We therefore could not directly evaluate the adaptive scenario.

However, it is unlikely that the almost ubiquitous pattern among invertebrates of larger eggs at lower temperatures could be a result of detailed fitness tradeoffs in such diverse organisms as butterflies, marine gastropods, and soil nematodes. Reduced cell size at higher temperatures has been demonstrated in fish blood cells, nematode cells, and various *Drosophila* epidermal cells (Van Voorhies, 1996; Azevedo, French & Partridge, 2002). One possible explanation for this general effect of temperature is the fundamental scaling relationships between oxygen supply and consumption. This predicts that cells or eggs produced at high temperatures should be smaller than those produced at lower temperatures (Woods, 1999). Unfortunately, we cannot evaluate this hypothesis without additional data. It would be necessary to measure oxygen gradients within the eggs themselves and demonstrate that levels are low enough in the centre of the egg to effect development (Woods, 1999).

To our knowledge, such data are not currently available for any marine invertebrate.

#### FEMALE EFFECT

The results obtained in the present study show that more than half of the variation in egg size and hatching size for each species of *Crepidula* was a result of differences among females. Such differences in egg size between females could result from genetic or environmental factors. Traits that are presumably highly linked to fitness and are therefore under selection are not expected to show high levels of genetic variation, although high heritabilities in egg size have been demonstrated for *Hydroides elegans* (Miles, Hadfield & Wayne 2007). Significant differences between females are frequently found in studies of marine invertebrate reproduction and most studies seek nongenetic factors to explain this variation. For example, in 13 species of sea urchins, significant differences were found between females whose eggs were measured on the same day (Lessios, 1987). Similarly, 20% of the variation in egg volume and 86% of the variation in egg energy content from mussels collected from several locations was a result of variation among females (Phillips, 2007). In these studies, as well as our own, it was not clear what differences among the females could contribute to such variation in egg size. Phillips (2007) ruled out effects of female body size and condition, and we found little indication of these factors having large effects in *Crepidula*. Among-female variation could be inflated by the artificial induction of spawning (Phillips, 2007), although this cannot be the case in *Crepidula*, which were allowed to spawn naturally. Similarly, the idea that small-scale differences in habitat could result in among-female variation in egg size would only apply

to field-collected animals, and not those raised their entire lives under laboratory conditions.

One factor that has been shown to affect egg size and quality in a number of studies is brood order. A reduction in egg size with successive broods has been found in a number of marine invertebrates (Gallager & Mann, 1986; Qian & Chia, 1992; Jones *et al.*, 1996; Ito, 1997). Because we only measured eggs from a single brood from each female, our experimental design does not allow us to test for an effect of brood order.

#### POPULATION DIFFERENTIATION

We showed that genetic differentiation in mtCOI sequences between populations in *C. ustulatulina* is accompanied by differentiation in offspring size and shape. Such differentiation in life history characters is not evident in *C. atrasolea*, which also does not show differentiation in COI. This is the first time that differences in population genetic structure among congeners have been linked to among-population differences in developmental features.

Inter-population differentiation in developmental features has been reported for barnacles (Barnes & Barnes, 1965), polychaetes (Bridges & Heppell, 1996), and fishes (Johnston & Leggett, 2002; Kokita, 2003), although, in most cases, there is no information on the genetic structure of the populations. This makes it difficult to determine whether such differences are due to genetic differentiation or plastic responses to environmental differences between the sites. A single previous study, using *Adalaria proxima*, demonstrated that animals from populations that were differentiated with respect to allozymes (Todd *et al.* 1998) also differed in egg size, brood size, and hatching size (Jones *et al.*, 1996). If such genetic differences in offspring size and other reproductive features are common among genetically distinct populations, then the starting conditions required for evolutionary divergence in mode of development may be more common than previously appreciated.

#### CONCLUSIONS

*Crepidula atrasolea* and *C. ustulatulina* raised under controlled conditions produce eggs with a wide range of diameters. In *C. atrasolea*, the average egg from the brood with the largest eggs was twice the volume of the average egg from the brood with smallest eggs; in *C. ustulatulina*, there was a three-fold difference in volume between the broods with the largest and smallest egg sizes. Documentation of such significant variation lays the foundation for the in-depth study of variation in offspring size, which is necessary to develop a comprehensive theory of the evolution of

offspring size (Bernardo, 1996). Bernardo advocates an approach that, instead of expecting a single optimal offspring size, embraces intraspecific variation and the complex interplay between optimal offspring size from both the parents' and offsprings' perspective, at the same time as taking into account plasticities and maternal effects. Certainly, our simple experiment yielded evidence for wide variation in offspring size and supported the idea that several factors play a roll in this variation.

From a macroevolutionary perspective, detailed studies of intraspecific variation are vital to bring a fuller understanding of the role of developmental variation in speciation. The link between genetic and developmental differentiation among populations is particularly important for understanding the diversification of marine invertebrates, where cryptic sister species with similar ecology often differ in development. Inter-population divergence in development could play a part in driving speciation in these groups, existing inter-population differences in development could be dragged along with speciation driven by other features, or differences in development among sibling species could evolve after speciation. There are few available data with which to assess these different scenarios. The results obtained in the present study, demonstrating genetic differentiation in offspring size among populations, lend support to the first two of these scenarios. However, more studies of intraspecific variation are necessary to assess the generality of this pattern and further distinguish between possible scenarios.

#### ACKNOWLEDGEMENTS

We thank S. Reed (Smithsonian Marine Station in Fort Pierce) and L. K. Dixon (Mote Marine Laboratory) for providing long-term field temperature data. The Smithsonian Institution's Scholarly Studies program and Marine Science Network provided financial support for this project. M. F. Vinasco, R. Rissanen, M. Venegas, and J. Sartu assisted in animal maintenance and photography and D. Ingrao provided assistance in the field. R. Robertson, H. Lessios, K. McDonald, and J. McAlister commented on previous versions of this manuscript. This is SMS publication #803.

#### REFERENCES

- Abramoff MD, Magelhaes PJ, Ram SJ. 2004. Image processing with ImageJ. *Biophotonics International* **11**: 36–42.  
 Allen JD. 2008. Size-specific predation on marine invertebrate larvae. *Biological Bulletin* **214**: 42–49.  
 Allen JD, Zakas C, Podolsky RD. 2006. Effects of egg size reduction and larval feeding on juvenile quality for a species

- with facultative-feeding development. *Journal of Experimental Marine Biology and Ecology* **331**: 186–197.
- Azevedo RBR, French V, Partridge L. 2002.** Temperature modulates epidermal cell size in *Drosophila melanogaster*. *Journal of Insect Physiology* **48**: 231–237.
- Barnes H, Barnes M. 1965.** Egg size, nauplius, and their variations with local, geographical and specific factors in some common cirripedes. *Journal of Animal Ecology* **34**: 391–402.
- Bengtson DA, Barkman RC, Berry WJ. 1987.** Relationship between maternal size egg diameter time of spawning season temperature and length of hatch of Atlantic silver-side, *Menidia menidia*. *Journal of Fish Biology* **31**: 697–704.
- Bernardo J. 1996.** The particular maternal effect of propagule size, especially egg size: patterns, models, quality of evidence and interpretations. *American Zoologist* **36**: 216–236.
- Bertram DF, Strathmann RR. 1998.** Effects of maternal and larval nutrition on growth and form of planktotrophic larvae. *Ecology* **79**: 315–327.
- Bridges TS, Heppell S. 1996.** Fitness consequences of maternal effects in *Streblospio benedicti* (Annelida: Polychaeta). *American Zoologist* **36**: 132–146.
- Chaparro OR, Oyarzun RF, Vergara AM, Thompson RJ. 1999.** Energy investment in nurse eggs and egg capsules in *Crepidula dilatata* Lamark (Gastropoda, Calyptraeidae) and its influence on hatching size of juveniles. *Journal of Experimental Marine Biology and Ecology* **232**: 261–274.
- Collin R. 2000.** Phylogeny of the *Crepidula plana* (Gastropoda: Calyptraeidae) cryptic species complex in North America. *Canadian Journal Zoology* **78**: 1500–1514.
- Collin R. 2001.** The effects of mode of development on phylogeography and population structure of North Atlantic *Crepidula* (Gastropoda: Calyptraeidae). *Molecular Ecology* **10**: 2249–2262.
- Collin R. 2002.** Another last word on *Crepidula convexa* and a description of *C. ustulatulina* sp. nov. (Gastropoda: Calyptraeidae) from the Gulf of Mexico. *Bulletin of Marine Science* **70**: 177–184.
- Collin R. 2003.** World-wide patterns of development in calyptraeid gastropods. *Marine Ecology Progress Series* **247**: 103–122.
- Collin R. 2004.** The loss of complex characters, phylogenetic effects, and the evolution of development in a family of marine gastropods. *Evolution* **58**: 1488–1502.
- Collin R, Chaparro OR, Winkler F, Veliz D. 2007.** Molecular phylogenetic and embryological evidence that feeding larvae have been reacquired in a marine gastropod. *Biological Bulletin* **212**: 83–92.
- Emlet RB. 1995.** Developmental mode and species geographic range in regular sea urchins (Echinodermata: Echinoidea). *Evolution* **49**: 476–489.
- Ernsting G, Isaaks A. 2000.** Ectotherms, temperature, and trade-offs: size and number of eggs in a carabid beetle. *American Naturalist* **155**: 804–813.
- Fischer K, Brakefield PM, Zwaan BJ. 2003.** Plasticity in butterfly egg size: why larger offspring at lower temperatures? *Ecology* **84**: 3138–3147.
- Fox CW. 1993.** The influence of maternal age and mating frequency on egg size and offspring performance in *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Oecologia* **96**: 139–146.
- Gallager SM, Mann R. 1986.** Growth and survival of larvae of *Mercenaria mercenaria* (L.) and *Crassostrea virginica* (Gmelin) relative to broodstock conditioning and lipid content of eggs. *Aquaculture* **56**: 105–121.
- George SB. 1996.** Echinoderm egg and larval quality as a function of adult nutritional state. *Oceanologia Acta* **19**: 297–308.
- George SB, Cellario C, Fenaux L. 1990.** Population differences in egg quality of *Arbacia lixula* (Echinodermata: Echinoidea): proximate composition of egg and larval development. *Journal of Experimental Marine Biology and Ecology* **141**: 107–118.
- Hadfield MG. 1989.** Latitudinal effects on juvenile size and fecundity in *Petalococonchus* (Gastropoda). *Bulletin of Marine Science* **45**: 369–376.
- Hadfield MG, Strathmann MF. 1996.** Variability, flexibility and plasticity in life histories of marine invertebrates. *Oceanologia Acta* **19**: 323–334.
- Hart MW. 1995.** What are the costs of small eggs size for a marine invertebrate with feeding planktonic larvae? *American Naturalist* **146**: 415–426.
- Hoagland KE. 1984.** Use of molecular genetics to distinguish species of the gastropod genus *Crepidula* (prosobranchia: Calyptraeidae). *Malacologia* **25**: 607–628.
- Hoagland KE. 1986.** Patterns of encapsulation and brooding in the Calyptraeidae (Prosobranchia: Mesogastropoda). *American Malacological Bulletin* **4**: 173–183.
- Honkoop PJC, Van Der Meer J. 1998.** Experimentally induced effects of water temperature and immersion time on reproductive output of bivalves in the Wadden Sea. *Journal of Experimental Marine Biology and Ecology* **220**: 227–246.
- Ito K. 1997.** Egg-size and -number variations related to maternal size and age, and the relationship between egg size and larval characteristics in and annual marine gastropod, *Halio japonica* (Opisthobranchia; Cephalaspidea). *Marine Ecology Progress Series* **152**: 187–195.
- Johnston TA, Leggett WC. 2002.** Maternal and environmental gradients in the egg size of an iteroparous fish. *Ecology* **83**: 1777–1791.
- Jones HL, Todd CD, Lambert WJ. 1996.** Intraspecific variation in embryonic and larval traits of the dorid nudibranch mollusc *Adalaria proxima* (Alder and Hancock) around the northern coasts of the British Isles. *Journal of Experimental Marine Biology and Ecology* **202**: 29–47.
- Kohn AJ, Perron FE. 1994.** *Life history and biogeography patterns in Conus*. Oxford Biogeography Series No. 8. Oxford: Clarendon Press, 106.
- Kokita T. 2003.** Potential latitudinal variation in egg size and number of a geographically widespread reef fish, revealed by common-environment experiments. *Marine Biology* **143**: 593–601.
- Lardies MA, Castilla JC. 2001.** Latitudinal variation in the reproductive biology of the commensal crab *Pinnaxodes*

- chilensis* (Decapoda: Pinnotheridae) along the Chilean coast. *Marine Biology* **139**: 1125–1133.
- Lessios HA. 1987.** Temporal and spatial variation in egg size of 13 Panamanian echinoids. *Journal of Experimental Marine Biology and Ecology* **114**: 217–240.
- Levitan DR. 2006.** Relationship between egg size and fertilization success in broadcast spawning marine invertebrates. *Integrative and Comparative Biology* **46**: 298–311.
- Marshall DJ, Keough MJ. 2003.** Effects of settler size and density on early post-settlement survival of *Ciona intestinalis* in the field. *Marine Ecology Progress Series* **259**: 139–144.
- Marshall DJ, Keough MJ. 2008.** The evolutionary ecology of offspring size in marine invertebrates. *Advances in Marine Biology* **53**: 1–60.
- Marshall DJ, Styan CA, Keough MJ. 2000.** Intraspecific co-variation between egg and body size affects fertilisation kinetics of free-spawning marine invertebrates. *Marine Ecology Progress Series* **195**: 305–309.
- McEdward LR, Carson SF. 1987.** Variation in organic content and its relationship with egg size in the starfish *Solaster stimpsoni*. *Marine Ecology Progress Series* **37**: 159–169.
- McEdward LR, Coulter LK. 1987.** Egg volume and energetic content are not correlated among sibling offspring of starfish: implications for life-history theory. *Evolution* **41**: 914–917.
- Miles CM, Hadfield MG, Wayne ML. 2007.** Heritability for egg size in the serpulid polychaete *Hydroides elegans*. *Marine Ecology Progress Series* **340**: 155–162.
- Moran AL, Emler RB. 2001.** Offspring size and performance in variable environments: field studies on a marine snail. *Ecology* **82**: 1597–1612.
- Moran AL, McAlister JS. 2009.** Egg size as a life history character of marine invertebrates: is it all it's cracked up to be? *Biological Bulletin* **216**: 226–242.
- Phillips NE. 2007.** Variability in egg size and energetic content among intertidal mussels. *Biological Bulletin* **212**: 12–19.
- Qian P-Y. 1994.** Effect of food quantity on growth and reproductive characteristics of *Capitella* sp. (Annelida: Polychaeta). *Invertebrate Reproduction and Development* **26**: 175–185.
- Qian P-Y, Chia F-S. 1992.** Effect of aging on reproduction in a marine polychaete *Capitella* sp. *Journal of Experimental Marine Biology and Ecology* **156**: 23–38.
- Quinn GP, Keough MJ. 2002.** *Experimental design and data analysis for biologists*. Cambridge: Cambridge University Press.
- Rivest BR. 1983.** Development and the influence of nurse egg allotment on hatching size in *Searlesia dira* (Reeve, 1846) (Prosobranchia: Buccinidae). *Journal of Experimental Marine Biology and Ecology* **69**: 217–241.
- Skadsheim A. 1989.** Regional variation in amphipod life history: effects of temperature and salinity on breeding. *Journal of Experimental Marine Biology and Ecology* **127**: 25–42.
- Todd CD, Lambert MJ, Thorpe JP. 1998.** The genetic structure of intertidal populations of two species of nudibranch molluscs with planktotrophic and pelagic lecithotrophic larval stages: are pelagic larvae 'for' dispersal? *Journal of Experimental Marine Biology and Ecology* **228**: 1–28.
- Van Voorhies WA. 1996.** Bergmann size clines: a simple explanation for their occurrence in ectotherms. *Evolution* **50**: 1259–1264.
- Woods HA. 1999.** Egg-mass size and cell size: effects of temperature on oxygen distribution. *American Zoologist* **39**: 244–252.